



GCSE
COMBINED SCIENCE: TRILOGY
8464/C/1H

Chemistry Paper 1H

Mark scheme

June 2022

Version: 1.0 Final Mark Scheme



2 2 6 G 8 4 6 4 C 1 H / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Copyright information

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2022 AQA and its licensors. All rights reserved.

Information to Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the examiner make their judgement
- the Assessment Objectives and specification content that each question is intended to cover.

The extra information is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent (for example, a scientifically correct answer that could not reasonably be expected from a student's knowledge of the specification).

2. Emboldening and underlining

- 2.1** In a list of acceptable answers where more than one mark is available 'any **two** from' is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **or**.
Alternative words in the mark scheme are shown by a solidus eg allow smooth / free movement.
- 2.4** Any wording that is underlined is essential for the marking point to be awarded.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of errors / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (indicated as * in example 1) are not penalised.

Example 1: What is the pH of an acidic solution?

[1 mark]

Student	Response	Marks awarded
1	green, 5	0
2	red*, 5	1
3	red*, 8	0

Example 2: Name **two** magnetic materials.

[2 marks]

Student	Response	Marks awarded
1	iron, steel, tin	1
2	cobalt, nickel, nail*	2

3.2 Use of symbols / formulae

If a student writes a chemical symbol / formula instead of a required chemical name, or uses symbols to denote quantities in a physics equation, full credit can be given if the symbol / formula is correct and if, in the context of the question, such action is appropriate.

3.3 Marking procedure for calculations

Marks should be awarded for each stage of the calculation completed correctly, as students are instructed to show their working. At any point in a calculation students may omit steps from their working. If a subsequent step is given correctly, the relevant marks may be awarded.

Full marks are **not** awarded for a correct final answer from incorrect working.

3.4 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.5 Errors carried forward

An error can be carried forward from one question part to the next and is shown by the abbreviation 'ecf'.

Within an individual question part, an incorrect value in one step of a calculation does not prevent all of the subsequent marks being awarded.

3.6 Phonetic spelling

Marks should be awarded if spelling is not correct but the intention is clear, **unless** there is a possible confusion with another technical term.

3.7 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.8 Allow

In the mark scheme additional information, 'allow' is used to indicate creditworthy alternative answers.

3.9 Ignore

Ignore is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

3.10 Do not accept

Do **not** accept means that this is a wrong answer which, even if the correct answer is given as well, will still mean that the mark is not awarded.

3.11 Numbered answer lines

Numbered lines on the question paper are intended to support the student to give the correct number of responses. The answer should still be marked as a whole.

4. Level of response marking instructions

Extended response questions are marked on level of response mark schemes.

- Level of response mark schemes are broken down into levels, each of which has a descriptor.
- The descriptor for the level shows the average performance for the level.
- There are two marks in each level.

Before you apply the mark scheme to a student's answer, read through the answer and, if necessary, annotate it (as instructed) to show the qualities that are being looked for. You can then apply the mark scheme.

Step 1: Determine a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the descriptor for that level.

The descriptor for the level indicates the different qualities that might be seen in the student's answer for that level. If it meets the lowest level then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer. With practice and familiarity you will find that for better answers you will be able to quickly skip through the lower levels of the mark scheme.

When assigning a level you should look at the overall quality of the answer. Do **not** look to penalise small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level.

Use the variability of the response to help decide the mark within the level, ie if the response is predominantly level 2 with a small amount of level 3 material it would be placed in level 2 but be awarded a mark near the top of the level because of the level 3 content.

Step 2: Determine a mark

Once you have assigned a level you need to decide on the mark. The descriptors on how to allocate marks can help with this. The exemplar materials used during standardisation will help. There will be an answer in the standardising materials which will correspond with each level of the mark scheme. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the example to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the example.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.

Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other valid points. Students do not have to cover all of the points mentioned in the indicative content to reach the highest level of the mark scheme.

You should ignore any irrelevant points made. However, full marks can be awarded only if there are no incorrect statements that contradict a correct response.

An answer which contains nothing of relevance to the question must be awarded no marks.

Question 1

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.1	any three from: <ul style="list-style-type: none"> • green solid / powder • colourless solution • blue solution formed • copper carbonate disappears • fizzing / effervescence or bubbles (of gas) • stops fizzing • solid / powder left at the end or copper carbonate left at the end 	ignore green copper carbonate allow colour (of solution) changes allow solid disappears ignore gas allow fizzing slows down allow (container) gets hot or allow temperature increases	3	AO2 AO3 5.4.2.2 5.4.2.3 RPA8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.2	filtration or filter		1	AO1 5.4.2.3 RPA8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.3	7		1	AO1 5.4.2.3 5.4.2.4 RPA8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.4	neutralisation	allow exothermic	1	AO1 5.4.2.2 5.4.2.4 RPA8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.5	83 (g at 80 °C)	allow a value in range 82–84 (g at 80 °C)	1	AO2
	32 (g at 20 °C)	allow a value in range 32–33 (g at 20 °C)	1	AO2
	(83–32 =) 51 (g)	allow a correct calculation using incorrectly read values for mass at 80 °C and/or 20 °C	1	AO3 5.4.2.3

Total Question 1		9
-------------------------	--	----------

Question 2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.1	K ₂ SO ₄		1	AO2 5.1.1.1 5.4.3.4 RPA9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.2	(volume of hydrogen) 30 (cm ³) and (volume of oxygen) 15 (cm ³)		1	AO2 5.4.3.4 RPA9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.3	(because) the ratio of volume of hydrogen : oxygen is 2 : 1		1	AO3 5.4.3.4 RPA9
	(and this is the) same as the ratio of hydrogen (atoms) : oxygen (atoms) in (formula of) H ₂ O		1	
	OR			
	(because) the ratio of volume of hydrogen : oxygen is not 2 : 1 (1)	must relate to the volumes given in question 02.2		
	(and this is) different to the ratio of hydrogen (atoms) : oxygen (atoms) in (formula of) H ₂ O (1)			

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.4	$9 \pm 3 \text{ cm}^3$		1	AO2 5.3.1.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.5	(conversion) $\left(\frac{25}{1000} = \right) 0.025 \text{ (dm}^3\text{)}$		1	AO2 5.3.2.5 5.4.3.4
	(concentration =) $\frac{0.86}{0.025}$ $= 34.4 \text{ (g per dm}^3\text{)}$	allow correct use of incorrect / no conversion	1	
	OR (conversion) $\frac{1000}{25} \text{ (1)}$ $= 40 \text{ (1)}$ (40×0.86) $= 34.4 \text{ (g per dm}^3\text{) (1)}$	allow 34 (g per dm ³)	1	
OR (concentration =) $\frac{0.86}{25} \text{ (1)}$ $= 0.0344 \text{ (1)}$ (conversion) (0.0344×1000) $= 34.4 \text{ (g per dm}^3\text{) (1)}$	allow correct use of incorrect / no conversion allow 34 (g per dm ³)			

Total Question 2		8
-------------------------	--	----------

Question 3

Question	Answers	Mark	AO / Spec. Ref.
03	Level 3: The method would lead to the production of a valid outcome. All key steps are identified and logically sequenced.	5–6	AO3
	Level 2: The method would not necessarily lead to a valid outcome. Most steps are identified, but the method is not fully logically sequenced.	3–4	AO3
	Level 1: The method would not lead to a valid outcome. Some relevant steps are identified, but links are not made clear.	1–2	AO1
	No relevant content.	0	
	<p>Indicative Content</p> <ul style="list-style-type: none"> • measure volume of (hydrochloric) acid • into a suitable container eg polystyrene cup • measure the initial temperature (of hydrochloric acid) • with a thermometer • add stated mass of one metal • stir <ul style="list-style-type: none"> • measure the highest temperature reached of the solution or • measure temperature reached after a set time period • determine the temperature difference <ul style="list-style-type: none"> • repeat <ul style="list-style-type: none"> • repeat for each metal • with same mass • in same physical state (powder, lump, etc) • with the same volume and / or concentration of (hydrochloric) acid <ul style="list-style-type: none"> • use results to arrange metals in order of reactivity • most reactive metal has the largest temperature change <p>to access level 3 there must be an indication of how the temperature change is determined with the same mass of the 3 different metals reacted with the same volume of (hydrochloric) acid</p>		5.1.4.2 RPA2

Total Question 3		6
-------------------------	--	----------

Question 4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.1	halogens		1	AO1 5.1.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.2	all have 7 electrons in <u>outer</u> shell or all have 7 <u>outer</u> electrons	allow energy level for shell allow same number of <u>outer</u> electrons allow one electron required to complete the <u>outer</u> shell	1	AO1 5.1.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.3	Cl ₂ O ₇		1	AO2 5.2.1.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.4	y-axis scale correct from –100 to –250 °C bar correctly plotted at –220 °C	allow a tolerance of $\pm \frac{1}{2}$ a small square	1 1	AO2 5.1.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.5	(the) molecules increase in size going down the group	allow converse explanation in terms of decreasing melting point allow atoms increase in size going down the group allow increase in number of electron shells going down group	1	AO3
	(so the) forces between the molecules increase or (so the) intermolecular forces increase		1	AO1
	(so the) melting points increase going down the group or (so the) melting points increase with increasing relative atomic mass	allow (so) more energy is needed to separate the molecules	1	AO1 5.1.2.6 5.2.2.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.6	(s)		1	AO3 5.1.2.6 5.2.2.1 5.2.2.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.7	condensation	allow condensing ignore evaporating and boiling	1	AO1 5.2.2.1

Total Question 4		10
-------------------------	--	-----------

Question 5


Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.1	fullerene	allow (carbon) nanotube do not accept Buckminsterfullerene	1	AO1 5.2.3.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.2	any one from: <ul style="list-style-type: none"> • conducts heat • conducts electricity • very high length to diameter ratio 	allow large surface area to volume ratio allow high tensile strength allow can trap other molecules / atoms / ions	1	AO2 5.2.3.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.3	other metal atoms have different sizes to aluminium atoms		1	AO1 5.2.2.7
	(so) the layers of aluminium atoms are distorted		1	
	(so) the layers cannot slide	allow (so) the atoms cannot slide over each other	1	
	(which) makes the alloy harder	allow (which) makes the alloy stronger	1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.4	covalent bonds (between atoms) in the chain intermolecular forces between the chains covalent bonds are strong and intermolecular forces are weak		1 1 1	AO1 5.2.2.5
Total Question 5			9	

Question 6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.1	<p>one shared pair in overlap</p> <p>6 non-bonding electrons in outer shell of chlorine</p>	<p>allow any combination of circles, dots, crosses or e^{-}</p> <p>do not accept extra electron(s) on outer shell of hydrogen</p> <p>ignore any inner shell electrons</p> <p>an answer of</p>  <p>scores 2 marks</p>	<p>1</p> <p>1</p>	<p>AO1 5.1.2.6 5.2.1.4</p>
Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.2	completely ionises in aqueous solution	allow completely dissociates in aqueous solution	1	AO1 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.3	fizzing / effervescence or magnesium disappears		1	AO3 5.4.2.5
	at a greater rate with a strong acid	allow converse with weak acid allow for 2 marks strong acid has a greater temperature increase	1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.4	(pH) decreases	allow pH is lower	1	AO1
	by (a unit of) 2		1	AO2 5.4.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.5	(bonds broken = $(4 \times 413) + \text{C}=\text{C} + 431 =$ $2083 + \text{C}=\text{C}$)		1	AO2 5.5.1.3
	(bonds made = $(346 + 339 + (5 \times 413) =)$ 2750)		1	
	(energy released = bonds made – bonds broken =) $56 = 2750 - [2083 + \text{C}=\text{C}]$)	allow correct use of incorrect value(s) from step 1 and / or step 2	1	
	($\text{C}=\text{C}$) = 611 (kJ/mol)		1	
Total Question 6			11	

Question 7

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.1	electrolysis	allow liquid for molten	1	AO3 5.4.1.3 5.4.3.2 5.4.3.3
	of molten compound (of metal Y)		1	
	OR			
	displacement (1)			
	by heating with a more reactive metal			
	or			
	by heating with potassium / magnesium (1)			

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.2	$4 \text{ Na} + \text{TiCl}_4 \rightarrow 4 \text{ NaCl} + \text{Ti}$	allow multiples	2	AO2 5.1.1.1 5.3.1.1
		allow 1 mark for NaCl and Ti with incorrect / no balancing		

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.3	$\text{Na} \rightarrow \text{Na}^+ + \text{e}^-$	ignore state symbols	2	AO2 5.4.1.2 5.4.1.4
		allow multiples		
		allow 1 mark for $\text{Na} \rightarrow \text{Na}^+ + \text{e}^-$ with incorrect balancing		

Question	Answers	Extra information	Mark	AO / Spec. Ref.		
07.4	<p>method 1:</p> <p>(moles of Al = $\frac{108}{27}$ =) 4</p> <p>(moles CuCl₂ = $\frac{1210}{134.5}$ =) 8.996</p> <p>(identifying limiting reactant)</p> <table border="1" data-bbox="296 719 719 853"> <tr> <td>4 moles Al gives 6 moles Cu</td> </tr> <tr> <td>8.996 moles CuCl₂ gives 8.996 moles Cu</td> </tr> </table> <p>therefore aluminium is the limiting reactant</p> <p>(mass of Cu = 2 × 3 × 63.5) = 6 × 63.5</p> <p>= 381 (g)</p>	4 moles Al gives 6 moles Cu	8.996 moles CuCl ₂ gives 8.996 moles Cu	<p>allow 9</p> <p>allow correct use of an incorrectly calculated value(s) for moles of Al and / or CuCl₂</p> <p><u>must</u> follow on from MP3</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>	<p>AO2 5.3.2.2 5.3.2.4</p>
4 moles Al gives 6 moles Cu						
8.996 moles CuCl ₂ gives 8.996 moles Cu						

<p>07.4 cont.</p>	<p>method 2:</p> $2\text{Al} + 3\text{CuCl}_2 \rightarrow 3\text{Cu} + 2\text{AlCl}_3$ <p>(2×27) (3×134.5) (3×63.5)</p> <p>54 (g) 403.5 (g) 190.5 (g)</p> <p>(1) (1) (1)</p> <p>(so)</p> <p>108 g Al (reacts with 807 g CuCl₂) to produce 381 g Cu (1)</p> <p>(so) there is excess CuCl₂</p> <p>or</p> <p>807 g CuCl₂ is less than 1210 g CuCl₂ (1)</p> <p>therefore aluminium is limiting reactant (1)</p> <p>method 3:</p> <p>134.5 g CuCl₂ produces 63.5 g Cu (1)</p> <p>(mass conversion 1.21 kg CuCl₂ => 1210 (g) (1)</p> <p>1210 g CuCl₂ produces $\left(\frac{63.5}{134.5} \times 1210\right)$ => 571 g Cu (1)</p> <p>54 g Al produces 190.5 g Cu (1)</p> <p>108 g Al produces $\left(\frac{190.5}{54} \times 108\right)$ => 381 (g) (1)</p> <p>(therefore) aluminium is the limiting reactant (1)</p>	<p>allow correct use of an incorrect calculation of mass for Al / CuCl₂ / Cu</p> <p><u>must</u> follow on from MP4 / MP5</p> <p>allow correct use of an incorrect / no conversion of mass of CuCl₂</p> <p><u>must</u> follow on from MP3 and MP5</p>		
--------------------------	--	--	--	--

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.5	delocalised electrons	allow free electrons	1	AO1 5.2.2.8
	carry (electrical) charge through the metal / sodium	ignore throughout for through ignore current / electricity MP2 is dependent upon MP1	1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.6	(conducts electricity) when liquid / molten	allow (conducts electricity) when dissolved in water	1	AO1 5.2.2.3
	or (conducts electricity) in (aqueous) solution		1	
	(because) <u>ions</u> (ions) are free to move or (ions) allow charge to flow		1	

Total Question 7		17
-------------------------	--	-----------